

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/337890100>

Symmetric Functions of the K -Fibonacci and K -Lucas Numbers

Article in *Asian-European Journal of Mathematics* · January 2021

DOI: 10.1142/S1793557121500315

CITATION

1

READS

400

4 authors:



Ali Boussayoud
University of Jijel

60 PUBLICATIONS 266 CITATIONS

[SEE PROFILE](#)



Mohamed Kerada
University of Jijel

38 PUBLICATIONS 193 CITATIONS

[SEE PROFILE](#)



Serkan Araci
Hasan Kalyoncu University

242 PUBLICATIONS 1,664 CITATIONS

[SEE PROFILE](#)



Mehmet Acikgoz
Gaziantep University

252 PUBLICATIONS 1,415 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



q-analogues of some special polynomials [View project](#)



Sums of Powers of Consecutive Integers [View project](#)

Symmetric functions of the k -Fibonacci and k -Lucas numbers

Ali Boussayoud* and Mohamed Kerada†

*LMAM Laboratory and Department of Mathematics
Mohamed Seddik Ben Yahia University, Jijel, Algeria*

**aboussayoud@yahoo.fr*

†mkerada@yahoo.fr

Serkan Araci*

*Department of Economics, Faculty of Economics
Administrative and Social Sciences*

Hasan Kalyoncu University

TR-27410 Gaziantep, Turkey

mtsirken@hotmail.com; serkan.araci@hku.edu.tr

Mehmet Acikgoz

Department of Mathematics, Faculty of Arts and Science

University of Gaziantep, TR-27310 Gaziantep, Turkey

acikgoz@gantep.edu.tr

Communicated by H. M. Srivastava

Received November 21, 2017

Accepted December 5, 2019

Published March 11, 2020

In this paper, we introduce a new operator in order to derive some new symmetric properties of k -Fibonacci and k -Lucas numbers and Fibonacci polynomials. By making use of the new operator defined in this paper, we give some new generating functions for k -Fibonacci and Pell numbers and Fibonacci polynomials.

Keywords: k -Fibonacci numbers; k -Lucas numbers; generating functions; Fibonacci polynomials.

AMS Subject Classification: 05E05, 11B39

1. Introduction and Notations

Fibonacci numbers and their generalizations have many interesting properties and applications to almost every field of science and art (e.g. see [27]). Fibonacci

numbers F_n are defined by the recurrence relation

$$\begin{cases} F_0 = 1, & F_1 = 1, \\ F_{n+1} = F_n + F_{n-1}, & n \geq 1. \end{cases}$$

There exist a lot of properties about Fibonacci numbers. In particular, there is a beautiful combinatorial identity to Fibonacci numbers [27].

$$F_n = \sum_{i=0}^{\lfloor \frac{n-1}{2} \rfloor} \binom{n-i-1}{i}. \quad (1.1)$$

From (1.1), Filipponi [22] introduced the incomplete Fibonacci numbers $F_n(s)$ and the incomplete Lucas numbers $L_n(s)$. They are defined by

$$\begin{aligned} F_n(s) &= \sum_{j=0}^s \binom{n-j-1}{j}, \quad \left(0 \leq s \leq \left\lfloor \frac{n-1}{2} \right\rfloor; n = 0, 1, 2, \dots\right), \\ L_n(s) &= \sum_{j=0}^s \binom{n-j}{j}, \quad \left(0 \leq s \leq \left\lfloor \frac{n}{2} \right\rfloor; n = 0, 1, 2, \dots\right). \end{aligned}$$

In [17], Djordjevic gave the incomplete generalized Fibonacci and Lucas numbers. In [18], Djordjevic and Srivastava defined incomplete generalized Jacobsthal and Jacobsthal–Lucas numbers. In [16], the authors defined the incomplete Fibonacci and Lucas numbers. For the systematic work related to Fibonacci and related numbers, we refer the readers to see the references [29–32].

On the other hand, many kinds of generalizations of Fibonacci numbers have been presented in the literature. In particular, a generalization is the k -Fibonacci numbers. For any positive real number k , the k -Fibonacci sequence, say $(F_{n,k})_{n \in \mathbb{N}}$, is defined recurrently by

$$\begin{cases} F_{k,0} = 1, & F_{k,1} = 1, \\ F_{k,n+1} = kF_{k,n} + F_{k,n-1}, & n \geq 1. \end{cases}$$

In [24], k -Fibonacci numbers were found by studying the recursive application of two geometrical transformations used in the four-triangle longest-edge (4TLE) partition. These numbers have been studied in several papers; see [24, 25].

For any positive real number k , the k -Lucas sequence, say $(L_{n,k})_{n \in \mathbb{N}}$, is defined recurrently by

$$\begin{cases} L_{k,0} = 2, & L_{k,1} = k, \\ L_{k,n+1} = kL_{k,n} + L_{k,n-1}, & n \geq 1. \end{cases}$$

If $k = 2$, we have the classical Pell numbers appears: $P_0 = 0, P_1 = 1$ and $P_{n+1} = 2P_n + P_{n-1}$ for $n \geq 1$, cf. [3].

In this contribution, we shall define a new useful operator denoted by $\delta_{e_1 e_2}^{-k}$ for which we can formulate, extend and prove new results based on our previous ones [3, 5]. In order to determine new generating functions of the products of some

known numbers and polynomials, we combine between our indicated past techniques and these presented polishing approaches.

Let k and n be two positive integer and $\{a_1, a_2, \dots, a_n\}$ are set of given variables, recall [20] that the k th elementary symmetric function $e_k(a_1, a_2, \dots, a_n)$ and the k th complete homogeneous symmetric function $h_k(a_1, a_2, \dots, a_n)$ are defined respectively by

$$e_k(a_1, a_2, \dots, a_n) = \sum_{i_1 + i_2 + \dots + i_n = k} a_1^{i_1} a_2^{i_2} \cdots a_n^{i_n}, \quad 0 \leq k \leq n,$$

with $i_1, i_2, \dots, i_n = 0$ or 1 ,

$$h_k(a_1, a_2, \dots, a_n) = \sum_{i_1 + i_2 + \dots + i_n = k} a_1^{i_1} a_2^{i_2} \cdots a_n^{i_n}, \quad 0 \leq k \leq n,$$

with $i_1, i_2, \dots, i_n \geq 0$.

First, we set $e_0(a_1, a_2, \dots, a_n) = 1$ and $h_0(a_1, a_2, \dots, a_n) = 1$ (by convention). For $k > n$ or $k < 0$, we set $e_k(a_1, a_2, \dots, a_n) = 0$ and $h_k(a_1, a_2, \dots, a_n) = 0$.

Definition 1.1. Let $E = \{e_1, e_2\}$ an alphabet, we define the symmetric function S_n associated with the alphabet E by

$$S_n(e_1 + e_2) = h_n(e_1, e_2) = \frac{e_1^{n+1} - e_2^{n+1}}{e_1 - e_2}, \quad (1.2)$$

with

$$\begin{aligned} S_0(e_1 + e_2) &= h_0(e_1, e_2) = 1, \\ S_1(e_1 + e_2) &= h_1(e_1, e_2) = e_1 + e_2, \\ S_2(e_1 + e_2) &= h_2(e_1, e_2) = e_1^2 + e_1 e_2 + e_2^2, \\ &\vdots \end{aligned}$$

Definition 1.2. Let A and B be any two alphabets, then we give $S_n(A - B)$ by the following form:

$$\frac{\prod_{b \in B} (1 - bz)}{\prod_{a \in A} (1 - az)} = \sum_{n=0}^{\infty} S_n(A - B) z^n \quad (1.3)$$

with the condition $S_n(A - B) = 0$ for $n < 0$ (see [1]).

Corollary 1.1. Taking $A = 0$ in (1.3) gives

$$\prod_{b \in B} (1 - bz) = \sum_{n=0}^{\infty} S_n(-B) z^n \quad (1.4)$$

Further, in the case $A = 0$ or $B = 0$, we have

$$\sum_{n=0}^{\infty} S_n(A - B) z^n = \sum_{n=0}^{\infty} S_n(A) z^n \sum_{n=0}^{\infty} S_n(-B) z^n.$$

Definition 1.3. Let g be any function on \mathbb{R}^n , then we consider the divided difference operator as the following form

$$\partial_{x_i x_{i+1}}(g) = \frac{g(x_1, \dots, x_i, x_{i+1}, \dots, x_n) - g(x_1, \dots, x_{i-1}, x_{i+1}, x_i, x_{i+2}, \dots, x_n)}{x_i - x_{i+1}}$$

(see [19]).

Definition 1.4 ([9]). The symmetrizing operator $\delta_{e_1 e_2}^k$ is defined by

$$\delta_{e_1 e_2}^k(f) = \frac{e_1^k f(e_1) - e_2^k f(e_2)}{e_1 - e_2} \quad (k \in \mathbb{N}). \quad (1.5)$$

Remark 1.1. Let $E = \{e_1, e_2\}$ an alphabet, we have

$$h_k(e_1, e_2) = S_k(e_1 + e_2) = \delta_{e_1 e_2}^k(e_1).$$

2. The Fibonacci Polynomials

Note that if k is a real variable x , then $F_{k,n} = F_{x,n}$ and they correspond to the Fibonacci polynomials defined by [24]

$$F_{n+1}(x) = \begin{cases} 1 & \text{if } n = 0, \\ x & \text{if } n = 1, \\ xF_n(x) + F_{n-1}(x) & \text{if } n \geq 2. \end{cases} \quad (2.1)$$

from where the first Fibonacci polynomials are

$$\begin{aligned} F_1(x) &= 1, \\ F_2(x) &= x, \\ F_3(x) &= x^2 + 1, \\ F_4(x) &= x^3 + 2x, \\ F_5(x) &= x^4 + 3x^2 + 1, \\ F_6(x) &= x^5 + 4x^3 + 3x, \\ F_7(x) &= x^6 + 5x^4 + 6x^2 + 1, \\ F_8(x) &= x^7 + 6x^5 + 10x^3 + 4x. \end{aligned}$$

And from these expression, as for the k -Fibonacci numbers, we can write [24]:

$$F_{n+1}(x) = \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n-i}{i} x^{n-2i} \quad (n \geq 0).$$

Note that $F_{2n}(0) = 0$ and $x = 0$ is the only real root, while $F_{2n+1}(0) = 1$ with no real roots. Also for $x = k \in \mathbb{N}$, we obtain the elements of the k -Fibonacci numbers.

By iterating recurrence relation of formula (2.1), the following property is straightforwardly deduced.

Proposition 2.1 ([26]). *For $1 \leq r \leq n-1$ holds:*

$$F_{n+1}(x) = F_r(x)F_{n-(r-2)}(x) + F_{r-1}(x)F_{n-(r-1)}(x).$$

Proposition 2.2 (Binet's formula). *The n th Fibonacci polynomial may be written as*

$$F_n(x) = \frac{\sigma^n - (-\sigma)^{-n}}{\sigma + \sigma^{-1}}, \quad \text{being } \sigma = \frac{x + \sqrt{x^2 + 4}}{2}. \quad (2.2)$$

Proof. Note that the characteristic equation for k -Fibonacci polynomials is $r^2 - x$. $r - 1 = 0$ with roots $r_1 = \sigma = \frac{x + \sqrt{x^2 + 4}}{2}$, and $r_2 = -\sigma^{-1}$, from where Formula (2.2) is deduced. \square

Proposition 2.3 ([25]). *(Asymptotic behavior of the quotient of consecutive terms).*

$$\text{If } \sigma = \frac{x + \sqrt{x^2 + 4}}{2}, \text{ then } \lim_{n \rightarrow \infty} \frac{F_{n+1}(x)}{F_n(x)} = \sigma.$$

As a consequence, the quotient between two consecutive terms of the k -Fibonacci numbers $\{F_{k;n}\} = \{0, 1, k, k^2 + 1, k^3 + 2k, \dots\}$ tends to the positive characteristic root $\sigma = \sigma_k$. For each integer k , $\sigma + \sigma_k$ is called the k th metallic ration [28]: Golden Ratio, for $k = 2$, and Bronze Ratio for $k = 3$.

Proposition 2.4 ([24]). *(Honsberger's formula). for n, m integers it holds:*

$$F_{m+n}(x) = F_{m-1}(x)F_n(x) + F_m(x)F_{n-1}(x).$$

3. On the Symmetric Functions of Some k -Fibonacci Numbers and Fibonacci Polynomials

In this part, we are now in a position to provide Theorem 3.1. Also, we derive the new generating functions of the products of some known numbers and polynomials.

Definition 3.1. The symmetrizing operator $\delta_{e_1 e_2}^{-k}$ is defined by [7]

$$\delta_{e_1 e_2}^{-k}(f) = \frac{e_2^k f(e_1) - e_1^k f(e_2)}{(e_1 e_2)^k (e_1 - e_2)} \quad (k \in \mathbb{N}).$$

Lemma 3.1 ([7]). *Let $E = \{e_1, e_2\}$, we define the operator $\delta_{e_1 e_2}^{-k}$ as follows:*

$$\delta_{e_1 e_2}^{-k} f(e_1) = \frac{-h_{k-1}(e_1, e_2)}{e_1^k e_2^k} f(e_1) + \frac{e_1^k}{e_1^k e_2^k} \partial_{e_1 e_2} f(e_1).$$

Theorem 3.1. Let E and A be two alphabets, respectively, $\{e_1, e_2\}$ and $\{a_1, a_2, \dots\}$, then we have

$$\begin{aligned} & \frac{\sum_{n=0}^{\infty} e_n(a_1, a_2, \dots, a_n) \delta_{e_1 e_2}^{k+n-1}(e_1) z^n}{\prod_{a \in A} (1 - a e_1 z) \prod_{a \in A} (1 - a e_2 z)} \\ &= \sum_{n=0}^{k-1} h_n(a_1, a_2, \dots, a_n) e_1^n e_2^n \delta_{e_1 e_2}^{k-n-1}(e_1) z^n \\ & \quad - e_1^k e_2^k z^{k+1} \sum_{n=0}^{\infty} h_{n+k+1}(a_1, a_2, \dots, a_n) \delta_{e_1 e_2}^n(e_1) z^n \end{aligned} \quad (3.1)$$

for all $k \in \mathbb{N}$.

Proof. By applying the operator $\delta_{e_1 e_2}^{-k}$ to the series $f(e_1) = (\prod_{a \in A} (1 - a e_1 z))^{-1}$, we have

$$\begin{aligned} \delta_{e_1 e_2}^{-k} f(e_1) &= \delta_{e_1 e_2}^{-k} \left(\sum_{n=0}^{\infty} e_n(a_1, a_2, \dots, a_n) e_1^n z^n \right)^{-1} \\ &= \frac{\frac{e_2^k}{\sum_{n=0}^{\infty} e_n(a_1, a_2, \dots, a_n) e_1^n z^n} - \frac{e_1^k}{\sum_{n=0}^{\infty} e_n(a_1, a_2, \dots, a_n) e_2^n z^n}}{e_1^k e_2^k (e_1 - e_2)} \\ &= \frac{\sum_{n=0}^{\infty} e_n(a_1, a_2, \dots, a_n) e_2^{n+k} z^n - \sum_{n=0}^{\infty} e_n(a_1, a_2, \dots, a_n) e_1^{n+k} z^n}{e_1^k e_2^k (e_1 - e_2) (\prod_{a \in A} (1 - a e_1 z)) (\prod_{a \in A} (1 - a e_2 z))} \\ &= \frac{-1}{e_1^k e_2^k} \left(\frac{\sum_{n=0}^{\infty} e_n(a_1, a_2, \dots, a_n) \delta_{e_1 e_2}^{k+n-1}(e_1) z^n}{\prod_{a \in A} (1 - a e_1 z) \prod_{a \in A} (1 - a e_2 z)} \right). \end{aligned}$$

On the other hand,

$$\begin{aligned} \delta_{e_1 e_2}^{-k} f(e_1) &= \delta_{e_1 e_2}^{-k} \left(\sum_{n=0}^{\infty} h_n(a_1, a_2, \dots, a_n) e_1^n z^n \right) \\ &= \frac{e_2^k \sum_{n=0}^{\infty} h_n(a_1, a_2, \dots, a_n) e_1^n z^n - e_1^k \sum_{n=0}^{\infty} h_n(a_1, a_2, \dots, a_n) e_2^n z^n}{e_1^k e_2^k (e_1 - e_2)} \\ &= \frac{1}{e_1^k e_2^k} \left(\sum_{n=0}^{\infty} h_n(a_1, a_2, \dots, a_n) \frac{e_2^k e_1^n - e_1^k e_2^n}{e_1 - e_2} z^n \right) \\ &= \frac{1}{e_1^k e_2^k} \left(\sum_{n=0}^{k-1} h_n(a_1, a_2, \dots, a_n) \frac{e_2^k e_1^n - e_1^k e_2^n}{e_1 - e_2} z^n \right. \\ & \quad \left. + \sum_{n=k+1}^{\infty} h_n(a_1, a_2, \dots, a_n) \frac{e_2^k e_1^n - e_1^k e_2^n}{e_1 - e_2} z^n \right) \end{aligned}$$

$$\begin{aligned}
&= \frac{-1}{e_1^k e_2^k} \left(\sum_{n=0}^{k-1} h_n(a_1, a_2, \dots, a_n) e_1^n e_2^n \delta_{e_1 e_2}^{k-n-1}(e_1) z^n - e_1^k e_2^k z^{k+1} \right. \\
&\quad \left. \times \sum_{n=0}^{\infty} h_{n+k+1}(a_1, a_2, \dots, a_n) \delta_{e_1 e_2}^n(e_1) z^n \right).
\end{aligned}$$

This completes the proof. \square

We now derive the new generating functions of the products of some known polynomials. Indeed, we consider Theorem 1 in order to derive k -Fibonacci numbers and Fibonacci polynomials if $k = 1$.

Theorem 3.2. *Let E and A be two alphabets, respectively, $\{e_1, e_2\}$ and $\{a_1, a_2\}$, then we have*

$$\begin{aligned}
&\sum_{n=0}^{\infty} h_{n+2}(a_1, a_2) \delta_{e_1 e_2}^n(e_1) z^n \\
&= \frac{e_1 e_2 a_1^2 a_2^2 z^2 - a_1 a_2 h_1(e_1, e_2) h_1(a_1, a_2) z + (a_1 + a_2)^2 - a_1 a_2}{\prod_{a \in A} (1 - a e_1 z) \prod_{a \in A} (1 - a e_2 z)}. \quad (3.2)
\end{aligned}$$

In the case $A = \{a_1\}$, based on Theorem 3.2, we deduce the following lemmas:

Lemma 3.2. *Given two alphabets $E = \{e_1, -e_2\}$ and $A = \{a_1\}$, we have*

$$\sum_{n=0}^{\infty} a_1^n \delta_{e_1[-e_2]}^n(e_1) z^n = \frac{1}{1 - a_1(e_1 - e_2)z - a_1^2 e_1 e_2 z^2}. \quad (3.3)$$

Lemma 3.3. *Given two alphabets $E = \{e_1, -e_2\}$ and $A = \{a_1\}$, we have*

$$\sum_{n=0}^{\infty} a_1^n \delta_{e_1[-e_2]}^{n+1}(e_1) z^n = \frac{e_1 - e_2 + e_1 e_2 z}{1 - a_1(e_1 - e_2)z - a_1^2 e_1 e_2 z^2}. \quad (3.4)$$

Assuming that $e_1 - e_2 = 1$, $e_1 e_2 = 1$ and $a_1 = 1$ in Eqs. (3.3) and (3.4), we obtain the generating functions given by Boussayoud *et al.* [2, 5] which represent:

- (1) The generating function of the Fibonacci numbers F_n .
- (2) The generating function of the Lucas numbers L_n .

Choosing e_1 and e_2 such that $\begin{smallmatrix} e_1 e_2 = 1 \\ e_1 - e_2 = k \end{smallmatrix}$ and substituting in (3.3) and (3.4) we end up with [3]

$$\sum_{n=0}^{\infty} \delta_{e_1[-e_2]}^n(e_1) z^n = \frac{1}{1 - k z - z^2} \quad (3.5)$$

and

$$\sum_{n=0}^{\infty} \delta_{e_1[-e_2]}^{n+1}(e_1) z^n = \frac{k+z}{1-kz-z^2}. \quad (3.6)$$

Thus, we deduce the following theorem.

Theorem 3.3 ([2]). *For $n \in \mathbb{N}$, the generating function of the k -Fibonacci numbers is given by*

$$\sum_{n=0}^{\infty} F_{k,n} z^n = \frac{1}{1-kz-z^2}.$$

Multiplying Eq. (3.5) by the variable $2+k^2$ and subtract it from (3.6) multiplying by the variable k yields

$$\sum_{n=0}^{\infty} [(2+k^2)\delta_{e_1[-e_2]}^n(e_1) - k\delta_{e_1[-e_2]}^{n+1}(e_1)] z^n = \frac{2-kz}{1-kz-z^2},$$

and we have the following theorem.

Theorem 3.4 ([2]). *For $n \in \mathbb{N}$, the generating function of the k -Lucas numbers is given by*

$$\sum_{n=0}^{\infty} L_{k,n} z^n = \frac{2-kz}{1-kz-z^2}. \quad (3.7)$$

- Put $k = 2$ in the relationship (3.7), we have

$$\sum_{n=0}^{\infty} Q_n t^n = \frac{2-2t}{1-2t-t^2},$$

which represents a generating function for Pell–Lucas numbers [5].

Choosing e_1 and e_2 such that $\begin{cases} e_1 e_2 = 1 \\ e_1 - e_2 = x \end{cases}$ and substituting in (3.3), we end up with

$$\sum_{n=0}^{\infty} \delta_{e_1[-e_2]}^n(e_1) z^n = \frac{1}{1-xz-z^2}, \quad \text{with } e_1 = \sigma = \frac{x + \sqrt{x^2+4}}{2}.$$

Thus, we get the following theorem.

Theorem 3.5. *We have the following a new generating function of the Fibonacci polynomials as*

$$\sum_{n=0}^{\infty} F_n(x) t^n = \frac{1}{1-xt-t^2}.$$

For the case $E = \{e_1, -e_2\}$ and $A = \{a_1, -a_2\}$ with replacing e_2 by $-e_2$, a_2 by $-a_2$ in (3.2), we have

$$\begin{aligned} & \sum_{n=0}^{\infty} h_{n+2}(a_1, [-a_2]) h_n(e_1, [-e_2]) z^n \\ &= \frac{-e_1 e_2 a_1^2 a_2^2 z^2 + a_1 a_2 h_1(e_1, [-e_2]) h_1(a_1, [-a_2]) z + (a_1 - a_2)^2 + a_1 a_2}{(1 - a_1 e_1 z)(1 + a_2 e_1 z)(1 + a_1 e_2 z)(1 - a_2 e_2 z)}. \end{aligned} \quad (3.8)$$

This case consists of three related parts. Firstly, the substitutions

$$\begin{cases} a_1 - a_2 = k, \\ a_1 a_2 = 1, \end{cases} \quad \text{and} \quad \begin{cases} e_1 - e_2 = x, \\ e_1 e_2 = 1, \end{cases}$$

in (3.8) give

$$\begin{aligned} & \sum_{n=0}^{\infty} h_{n+2}(a_1, [-a_2]) h_n(e_1, [-e_2]) z^n \\ &= \frac{k^2 + 1 + kxz - z^2}{1 - kxz - (x^2 + k^2 + 2)z^2 - kxz^3 + z^4} \\ &= \sum_{n=0}^{\infty} F_{k,n+2} F_n(x) z^n. \end{aligned}$$

From which we have the following theorem.

Theorem 3.6. *We have the following a new generating function of the product of k -Fibonacci numbers and Fibonacci polynomials as*

$$\sum_{n=0}^{\infty} F_{k,n+2} F_n(x) z^n = \frac{k^2 + 1 + kxz - z^2}{1 - kxz - (x^2 + k^2 + 2)z^2 - kxz^3 + z^4}. \quad (3.9)$$

- Put $k = 1$ in the relationship (3.9), we have

$$\sum_{n=0}^{\infty} F_{n+2} F_n(x) z^n = \frac{2 + xz - z^2}{1 - xz - (x^2 + 3)z^2 - xz^3 + z^4},$$

which represents a new generatings functions of the product of Fibonacci numbers and Fibonacci polynomials.

- Put $k = 2$ in the relationship (3.9), we have

$$\sum_{n=0}^{\infty} P_{n+2} F_n(x) z^n = \frac{5 + 2xz - z^2}{1 - 2xz - (x^2 + 6)z^2 - 2xz^3 + z^4},$$

which represents a new generatings functions of the product of Pell numbers and Fibonacci polynomials.

Secondly, by making the following restrictions:

$$\begin{cases} a_1 - a_2 = k, \\ a_1 a_2 = 1, \end{cases} \quad \text{and} \quad \begin{cases} e_1 - e_2 = k, \\ e_1 e_2 = 1, \end{cases}$$

in (3.8) gives

$$\begin{aligned} \sum_{n=0}^{\infty} h_{n+2}(a_1, [-a_2]) h_n(e_1, [-e_2]) z^n &= \frac{k^2 + 1 + k^2 z - z^2}{1 - k^2 z - 2(k^2 + 1)z^2 - k^2 z^3 + z^4} \\ &= \sum_{n=0}^{\infty} F_{k,n+2} F_{k,n} z^n \end{aligned} \quad (3.10)$$

representing a new generating function of k -Fibonacci numbers $F_{k,n}$.

On the other hand, we consider

$$\begin{aligned} \sum_{n=0}^{\infty} F_{k,n+2} F_{k,n} z^n &= \sum_{n=0}^{\infty} (F_{k,n+1} + F_{k,n}) F_{k,n} z^n \\ &= \sum_{n=0}^{\infty} F_{k,n+1} F_{k,n} z^n + \sum_{n=0}^{\infty} F_{k,n}^2 z^n. \end{aligned}$$

Since

$$\sum_{n=0}^{\infty} F_{k,n}^2 z^n = \frac{1 - z^2}{1 - k^2 z - 2(k^2 + 1)z^2 - k^2 z^3 + z^4} \quad (\text{see [3]})$$

we have

$$\begin{aligned} \sum_{n=0}^{\infty} F_{k,n+2} F_{k,n} z^n &= \frac{k^2(1 + z)}{1 - k^2 z - 2(k^2 + 1)z^2 - k^2 z^3 + z^4} \\ &\quad + \frac{1 - z^2}{1 - k^2 z - 2(k^2 + 1)z^2 - k^2 z^3 + z^4}, \end{aligned}$$

from those applications, we deduce the following theorem.

Theorem 3.7. *We have the following a new generating function of the product of two consecutive k -Fibonacci numbers as*

$$\sum_{n=0}^{\infty} F_{k,n+1} F_{k,n} z^n = \frac{k^2(1 + z)}{1 - k^2 z - 2(k^2 + 1)z^2 - k^2 z^3 + z^4}. \quad (3.11)$$

- Put $k = 1$ in the relationship (3.10) and (3.11), we obtain the following results.

Corollary 3.1. *We have the following a new generating function of the product of Fibonacci numbers as*

$$\sum_{n=0}^{\infty} F_{n+2} F_n z^n = \frac{2 + z - z^2}{1 - z - 4z^2 - z^3 + z^4}.$$

Corollary 3.2. For $n \in \mathbb{N}$, the generating function of the product of two consecutive Fibonacci numbers is given by

$$\sum_{n=0}^{\infty} F_{n+1} F_n z^n = \frac{1}{1 - 2z - 2z^2 + z^3}.$$

- Put $k = 2$ in the relationship (3.10) and (3.11), we obtain the following results.

Corollary 3.3. We have the following a new generating function of the product of Pell numbers as

$$\sum_{n=0}^{\infty} P_{n+2} P_n z^n = \frac{5 + 4z - z^2}{1 - 4z - 10z^2 - 4z^3 + z^4}.$$

Corollary 3.4. We have the following a new generating function of the product of two consecutive Pell numbers as

$$\sum_{n=0}^{\infty} P_{n+1} P_n z^n = \frac{4}{1 - 5z - 5z^2 + z^3}.$$

4. Conclusion

In this paper, we have derived new theorems in order to determine generating functions of k -Fibonacci numbers, k -Lucas numbers and Fibonacci polynomials. The derived theorems and lemmas are based on symmetric functions and products of these numbers and polynomials.

References

1. A. Abderrezzak, Généralisation de la transformation d'Euler d'une série formelle, *Adv. Math.* **103** (1994) 180–195.
2. A. Boussayoud, M. Boulyer and M. Kerada, On some identities and symmetric functions for Lucas and Pell numbers, *Electron. J. Math. Anal. Appl.* **5** (2017) 202–207.
3. A. Boussayoud and N. Harrouche, Complete symmetric functions and k -Fibonacci numbers, *Commun. Appl. Anal.* **20** (2016) 457–467.
4. A. Abderrezzak, M. Kerada and A. Boussayoud, Generalization of some hadamard product, *Commun. Appl. Anal.* **20** (2016) 301–306.
5. A. Boussayoud, M. Kerada and M. Boulyer, A simple and accurate method for determination of some generalized sequence of numbers, *Int. J. Pure Appl. Math.* **108** (2016) 503–511.
6. A. Boussayoud, A. Abderrezzak and M. Kerada, Some applications of symmetric functions, *Integers* **15** (2015) A#48, 1–7.
7. A. Boussayoud and R. Sahali, The application of the operator $L_{b_1 b_2}^{-k}$ in the series $\sum_{j=0}^{+\infty} a_j b_1^j z^j$, *J. Adv. Res. Appl. Math.* **7** (2015) 68–75.
8. A. Boussayoud, M. Kerada, R. Sahali and W. Rouibah, Some applications on generating functions, *J. Concr. Appl. Math.* **12** (2014) 321–330.
9. A. Boussayoud, M. Kerada and A. Abderrezzak, A generalization of some orthogonal polynomials, *Springer Proc. Math. Stat.* **41** (2013) 229–235.

10. A. T. Benjamin and J. J. Quinn, Proofs that really count: The art of combinatorial proof, (Mathematical Association of America, Washington, DC, 2003).
11. A. F. Horadam, Basic properties of a certain generalized sequence of numbers, *Fibonacci Q.* **3** (1965) 161–176.
12. A. F. Horadam, Generating functions for powers of a certain generalized sequence of numbers, *Duke Math. J.* **32** (1965) 437–446.
13. A. F. Horadam and J. M. Mahon, Pell and Pell-Lucas polynomials, *Fibonacci Q.* **23** (1985) 7–20.
14. A. Lascoux and A. M. Fua, Partition analysis and symmetrizing operators, *J. Comb. Theory A.* **109** (2005) 339–343.
15. C. Bolat and H. Kose, On the properties of k -Fibonacci numbers, *Int. J. Contemp. Math. Sci.* **5** (2010) 1097–1105.
16. D. Tasci and M. Cetin Firengiz, Incomplete Fibonacci and Lucas p -numbers, *Math. Comput. Model.* **52** (2010) 1763–1770.
17. G. B. Djordjevic, Generating functions of the incomplete generalized Fibonacci and generalized Lucas numbers, *Fibonacci Q.* **42** (2004) 106–113.
18. G. B. Djordjevic and H. M. Srivastava, Incomplete generalized Jacobsthal and Jacobsthal-Lucas numbers, *Math. Comput. Model.* **42** (2005) 1049–1056.
19. I. G. Macdonald, *Symmetric Functions and Hall polynomials* (Oxford University Press, 1979).
20. M. Merca, A Generalization of the symmetry between complete and elementary symmetric functions, *Indian J. Pure Appl. Math.* **45** (2014) 75–89.
21. P. A. Panzone, On the generating functions of Mersenne and Fermat primes, *Collect. Math.* **63** (2012) 59–69.
22. P. Filippini, Incomplete Fibonacci and Lucas numbers, *Rend. Circ. Mat. Palermo.* **45** (1996) 37–56.
23. S. Araci, Novel identities involving Genocchi numbers and polynomials arising from applications of umbral calculus, *Appl. Math. Comput.* **233** (2014) 599–607.
24. S. Falcon and A. Plaza, On the Fibonacci k -numbers, *Chaos Solitons Fractals* **32** (2007) 1615–1624.
25. S. Falcon and A. Plaza, The k -Fibonacci sequence and the Pascal 2-triangle, *Chaos, Solitons Fractals* **33** (2008) 38–49.
26. S. Falcon and A. Plaza, On k -Fibonacci sequences and polynomials and their derivatives, *Chaos, Solitons Fractals* **39** (2009) 1005–1019.
27. T. Koshy, *Fibonacci and Lucas Numbers with Applications* (Wiley-Interscience, 2001).
28. V. W. Spinadel, Themetallc means family and forbidden symmetries, *Int. Math. J.* **2** (2002) 279–288.
29. H. M. Srivastava, N. Tuglu and M Cetin, Some results on the q -analogues of the incomplete Fibonacci and Lucas polynomials, *Miskolc Math. Notes* **20** (2019) 511–524.
30. G. B. Djordjevic and H. M. Srivastava, Some generalizations of certain sequences associated with the Fibonacci numbers, *J. Indonesian Math. Soc.* **12** (2006) 99–112.
31. G. B. Djordjevic and H. M. Srivastava, Some generalizations of the incomplete Fibonacci and the incomplete Lucas polynomials, *Adv. Stud. Contemp. Math.* **11** (2005) 11–32.
32. R. K. Raina and H. M. Srivastava, A class of numbers associated with the Lucas numbers, *Math. Comput. Model.* **25**(7) (1997) 15–22.